

Opportunistic Infections Amongst HIV Infected Children and Correlation with Who Clinical Staging in Owerri, South East Nigeria

Chinelo Vivian Okeke ^{1,2}, Samson Dayo Ejikunle ², Bede Ikenna Nnolim ³, Idaboh Lilian Oghenetega ⁴, Ganiyat Omobola Yusuf ⁵, Maria-Lauretta Chito Orji ⁶,

¹Consultant Pediatrician, National Obstetric Fistula Center Abakaliki, Ebonyi State, South East Nigeria.

²Consultant Gynecologist, Imo State University, Teaching Hospital Orlu Imo State, South East Nigeria.

³Consultant Physician, Alex Ekueme Federal University, Teaching Hospital Abakaliki, Ebonyi State, South East Nigeria.

⁴Consultant Pediatrician, Federal Medical Center, Asaba Delta State, South Nigeria.

⁵Consultant Pediatrician, Federal Medical Center, Keffi Nassarawa State, North Central Nigeria.

⁶Consultant Pediatrician, Alex Ekueme Federal University, Teaching Hospital Abakaliki, Ebonyi State, South East Nigeria.

*Corresponding author: Okeke Chinelo Vivian; chinelookekemail@gmail.com

Abstract

Background: Opportunistic infections are the most common cause of death among children living with Human Immunodeficiency Virus/Acquired Immune Deficiency Syndrome. They take advantage of the weakened immune system and cause devastating illnesses. Studies with respect to the prevalence of opportunistic infections in HIV infected Nigerian children are very limited. This study aimed to determine the prevalence of opportunistic infections in HIV infected children at Federal University Teaching Hospital Owerri and to correlate it with World Health Organization clinical staging. **Methods:** 100 HIV infected children aged 2-16 years were recruited consecutively at the pediatric infectious disease clinic of the Federal University Teaching Hospital Owerri. Diagnostic criteria for opportunistic infections in HIV infected children were based on the hospital's standard clinical protocols aligned with national and international guidelines. They were classified into World Health Organization clinical staging. They were further categorized based on CD4 count values using World Health Organization classification of immune deficiency. **Result:** The mean age of participants was 9.73 years. Majority of participants (59%) were in World Health Organization Clinical Stage I and (1%) was in stage IV. Pneumonia (68%) was the most common opportunistic infection recorded among participants. Fifty seven percent of subjects had CD4 counts > 500cells/mm³ while 17% had CD4 counts < 200cells/mm³. There was significant association between opportunistic infections and WHO clinical Staging. Place of residence and socioeconomic status were found to be significant risk factors for development of opportunistic infections. **Conclusion:** In this study, the prevalence of Opportunistic infections in HIV infected children at Federal University Teaching Hospital Owerri remained high. Opportunistic Infections majorly occurred with worsening World Health Organisation clinical staging, low socioeconomic status and inhabitants of rural areas. We recommend consideration be given to early screening and treatment of Opportunistic Infections in HIV infected children.

Keywords: Opportunistic infections, HIV, Children, Immune suppression.

Background

The global Human Immunodeficiency Virus (HIV) and Acquired Immunodeficiency Syndrome (AIDS) epidemic have become one of the greatest threats to human health and development, particularly in less developed countries ^[1]. Nigeria has the second largest HIV epidemic following South Africa and the highest burden of pediatric HIV and AIDS in the world ^[2]. As at 2021, the Nigerian national prevalence of HIV infection is estimated at 1.4%. Twelve percent of people living with HIV infection are children

aged 0 to 14 years, while 8% are adolescents aged 10 to 19 years ^[3]. Profound immune suppression occurs in HIV infected children. The hallmark of this process is the depletion of CD4+ lymphocytes, which predisposes the patient to develop a variety of opportunistic infections ^[4].

Opportunistic infections (OIs) are the most common cause of death among children living with HIV/AIDS. They are called opportunistic because they take advantage of the weakened immune system and cause devastating illnesses. Common OIs in HIV infected children include Tuberculosis, Candida,

Pneumocystis jirovecii Pneumonia, chronic diarrhea, invasive and recurrent fungi, cryptosporidiosis, cytomegalovirus, varicella zoster virus, Herpes simplex virus and Toxoplasmosis. In children, OIs have a more aggressive course compared to adults [5]. The World Health Organization (WHO) recommends a range of medical interventions to reduce OIs. This includes chemoprophylaxis, immunization, reduction of exposure and early initiation of highly active anti-retroviral therapy (HAART) [6]. Failure of early treatment of OIs in people living with HIV (PLHIV) results to poor quality of life, impaired patient's response to anti-retroviral therapy, increased disease progression and increased medical cost [7,8].

Most reports on the magnitude of OIs in HIV infected children are from North America and Europe [9]. Ravichandra *et al* [10] reported a 56% prevalence of opportunistic infections in HIV infected children in India.

The burden of OIs in HIV infected children in low and middle-income countries including Nigeria are poorly documented. However, Imade PE and Eghafona NO [11] reported that malaria parasite infection has the highest prevalence among OIs in HIV infected children in Benin City Nigeria. This study therefore was undertaken to determine the prevalence of OIs in HIV infected children attending the pediatric infectious disease clinic at Federal University Teaching Hospital Owerri. Results obtained from this study will add to existing data on burden of OIs in HIV infected children. It will also serve as a reference for clinicians whom will partake in similar research.

Objectives

To determine the prevalence of opportunistic infections in HIV infected children at Federal University Teaching Hospital Owerri.

To correlate opportunistic infections with WHO clinical staging of HIV infection

To determine the socio demographic factors associated with opportunistic infections

Methods

Study setting

Owerri is the capital of Imo State in the South East Geopolitical Zone of Nigeria [12]. Imo state has an estimated population of about 4,927,563 with about 403,425 people living in Owerri [13]. Federal University teaching hospital Owerri is the major tertiary health care facility in the state which serves as a referral centre for those living in Owerri and the surrounding towns as well as neighbouring states of Abia, Rivers and Anambra.

Study design

A hospital based cross-sectional study.

Study period

Conducted from October 2018 to January 2019 at the paediatric infectious disease clinic of Federal University teaching hospital Owerri Imo State.

Inclusion criteria

All HIV positive children aged 2-16 years whose parents consented to the study.

Exclusion criteria

HIV positive children whose parents declined consent.

Study population

HIV positive children aged 2-16 years.

Sampling procedure

100 HIV positive children that met the inclusion criteria were enrolled consecutively.

Data collection

A structured interviewer administered questionnaire was developed and administered to parents and guardians, to obtain child's demographic data, mode of HIV transmission, current or past history of opportunistic infections, route of HIV transmission. Based on clinical presentations, the children were categorized into various WHO clinical staging. Diagnostic criteria for opportunistic infections in HIV infected children were based on the hospital's standard clinical protocols aligned with national and international guidelines. CD4 count was also determined using CY flow methods. Other investigations done include full blood count, sputum for AFB, chest x-ray, mantoux, blood culture, stool MCS, urine MCS, ultrasound scan of abdomen/chest, cerebrospinal fluid analysis and culture sensitivity. Investigations were done according to clinical presentation of patients.

Variables

The primary outcome of interest was prevalence of opportunistic infections in HIV infected children. The primary exposures and predictors included WHO clinical staging, low socio-economic status, place of inhabitant. These variables were selected based on previously published research works suggesting they influence the likelihood of opportunistic infections in HIV infected children. Diagnostic criteria for opportunistic infections in HIV infected children were based on the hospital's standard clinical protocols aligned with national and international guidelines. These criteria were used to classify the primary diagnosis and assess the presenting condition's severity.

Bias

Several strategies were employed to minimize bias in the study. Selection bias was addressed by applying strict inclusion and exclusion criteria, ensuring that the study population was representative of the broader pediatric population seen in the HIV clinic. Information bias was minimized through standardized data collection methods and the cross-verification of data sources.

Data analysis

The data collected was coded and imputed into Statistical Package for Social Sciences (SPSS) version 20.0. Frequency tables and figures were used to present relevant variables.

Quantitative variables were used to summarize descriptive statistics (mean and standard deviation, age and social class). Qualitative variables were summarized as proportions. The Chi-square test was used to compare associations between proportions.

Ethical approval

Ethical approval was obtained from the ethics committee of Federal Medical Center Owerri (FMC/OW/IIREC/182). Informed consent of the parents and assent from children 7 years and above were obtained for the study and confidentiality maintained.

Results

A total of 100 patients were recruited from October 2018 to January 2019. Fifty-one were females while 49 were males. Majority 44 (44%) are from middle socioeconomic class. Hundred percent of study population were Christians 100 (100%) and 65 (65%) are rural dwellers. Majority of participants (57) had CD4

cell count above 500 cells/mm³ and (17) have CD4 below 200cell/mm³ as shown in Table I.

Of the 100 subjects, 59 were in WHO clinical Stage 1 and only 1 was in stage 4. Subjects on ART were 93 while 7 were HAART naïve. It was observed that 20(63%) participants had Pneumonia, 5(16%) had oral thrush, 4(12%) had Pulmonary Tuberculosis and 3(9%) people had herpes zoster. There was a significant association between PTB, Pneumonia, Oral Thrush and WHO clinical Staging as depicted in Table II.

The WHO clinical staging of HIV and place of residence were found to be significant risk factors for development of opportunistic infections. Of all opportunistic infections, 71% (23) occurred among rural dwellers and this was statistically significant, (p-value 0.003). Similarly prevalence of opportunistic infections increased with worsening WHO stage and this was statistically significant (p-value 0.01). Prevalence of all the studied infections increased with lower social class. This was not statistically significant (p-value 0.92) as shown in Table III.

Table 1: Socio demographic Characteristics and CD4 levels of Participants

| Variable | Frequency | Percent (%) |
|---|------------------|----------------|
| Gender | | |
| Female | 51 | 51 |
| Male | 49 | 49 |
| Religion | | |
| Christianity | 100 | 100 |
| Social Class | | |
| Upper | 20 | 20 |
| Middle | 44 | 44 |
| Lower | 36 | 36 |
| Place of Residence | | |
| Rural | 65 | 65 |
| Urban | 35 | 35 |
| CD4 Cell level of Participants | Frequency | Percent |
| <200 | 17 | 17 |
| 200-349 | 13 | 13 |
| 350 -499 | 13 | 13 |
| >500 | 57 | 57 |
| Total | 100 | 100 |
| There were slightly more females than males, all participants were Christians, majority were from middle class, and lived in rural areas. Furthermore, 57% of study subjects have adequate immunity while 17% were immune suppressed. | | |

Table II: Who Staging, Use of HAART, Opportunistic infection and Correlation between opportunistic infections and WHO Staging

| | | |
|--|--------------------|-------------------|
| WHO Staging | | |
| Variables | Frequency | Percent |
| 1 | 59 | 59 |
| 2 | 16 | 16 |
| 3 | 24 | 24 |
| 4 | 1 | 1 |
| Total | 100 | 100 |
| Use of HAART | Frequency | Percent |
| Yes | 93 | 93 |
| No | 7 | 7 |
| Total | 100 | 100 |
| Opportunistic infection | Number | Percentage |
| Pneumonia | 20 | 63% |
| Oral thrush | 5 | 16% |
| PTB | 4 | 12% |
| Herpes zoster | 3 | 9% |
| Correlation between opportunistic infections and WHO Staging | | |
| Variables | WHO Staging | |
| | Rho | p-value |
| PTB | 0.30 | 0.003* |
| Herpes Zoster | 0.16 | 0.11 |
| Pneumonia | 0.73 | <0.001 |
| Oral thrush | 0.33 | 0.001* |
| Majority of study participants (51%) were in WHO stage one while only 1 patient was in WHO stage 4. 93% were on antiretroviral drugs while 7 patients were HAART naïve. 63% had pneumonia, the second most common opportunistic infection was oral thrush followed by pulmonary TB, herpes zoster was the least common. There was significant association between PTB, Herpes zoster, Pneumonia, oral thrush and WHO | | |

clinical staging of HIV.

rho Spearman Correlation coefficient**Table III: Association between Risk factors and Development of Opportunistic Infections**

| Variables | Opportunistic Infections n (%) | | | | Total | χ^2 [¥] | p-value |
|--------------------|--------------------------------|------------------|---------------------|------------------|-------------------|-----------------------|---------|
| Domestic Animals | Herpes Zoster | PTB | Bacterial Pneumonia | Oral Thrush | | | |
| Fowl/Ducks/Turkey | 1(50.0) | 1(100.0) | 8(88.9) | 1(100.0) | 11(84.6) | | |
| Goats/Sheep | 1(50.0) | 0 | 0 | 0 | 1 (7.7) | 4.88 | 0.55 |
| Dogs | 0 | 0 | 1(11.1) | 0 | 1 (7.7) | | |
| Total | 2(100.0) | 1(100.0) | 9(100.0) | 1(100.0) | 13(100.0) | | |
| Place of Residence | | | | | | | |
| Rural | 2 (66.6) | 4(100.0) | 14 (80.0) | 3 (60.0) | 23(71.9) | | |
| Urban | 1 (33.3) | 0 | 6 (20.0) | 2 (40.0) | 9 (28.1) | 14.08 | 0.003 |
| Total | 3 (100.0) | 4(100.0) | 20 (100.0) | 5 (100.0) | 32(100.0) | | |
| WHO Stage | | | | | | | |
| 1 | 0 | 0 | 0 | 0 | 0 | | |
| 2 | 2 (33.3) | 0 | 0 | 0 | 2 (6.3) | 11.14 | 0.01 |
| 3 | 1 (66.6) | 4 (100.0) | 20 (100.0) | 5(100.0) | 30 (93.7) | | |
| 4 | 0 | 0 | 0 | 0 | 0 | | |
| Total | 3 (100.0) | 4 (100.0) | 20 (100.0) | 5(100.0) | 32 (100.0) | | |
| Social Class | | | | | | | |
| Upper | 0 | 0 | 0 | 0 | 0 | | |
| Middle | 0 | 0 | 2 (14.3%) | 1 (25.0) | 3 (13.0) | 2.04 | 0.92 |
| Lower | 3 (100.0) | 2 (100.0) | 12 (85.7) | 3 (75.0) | 20 (87.0) | | |
| Total | 3 (100.0) | 2 (100.0) | 14(100.0) | 4 (100.0) | 23(100.0) | | |

Opportunistic infections were more common in those that lived in rural areas, low socio economic class, increasing WHO clinical staging and those whose parents reared fowls.

¥ Likelihood ratio, Table has many small variables (less than 5) as such odds ratio will be misleading

Discussion

The prevalence of opportunistic infections in HIV infected children aged 2-16years attending the HIV care/treatment center of Federal Teaching Hospital Owerri South East Nigeria was 32%. This finding is comparable to the 31.6% prevalence of OIs in HIV infected children in the study by Mamuru et al in Ethiopia^[14]. The prevalence of OIs in this study was also similar to the 28.8 incidence of OIs per 100 HIV infected children in an Asian study by Wasana Prasitsuebsai *et al.*^[15].

However, the prevalence of OIs in this study is lower than the 56% prevalence of OIs obtained in the Indian study by Ravichandra *et al.*^[10]. The difference in prevalence may be due to the fact that majority of patients in the Indian study were severely immune compromised, while majority of participants in this study were not severely immune compromised and were on HAART. Also, 98% of study participants washed hands regularly before a meal, 92% washed fruits regularly before eating and 75% drank water from bore hole. These simple acts of hygiene would have drastically reduced the risk of contracting infections.

Pneumonia was the most common opportunistic infection (63%) followed by oral thrush (16%). Tuberculosis and herpes zoster was seen in 12% and 9% of children respectively. Similarly, a study from Latin America,^[16] revealed that bacterial pneumonia is the most common OI. However, in contrast to our study, Imade E and Eghafona NO^[11] reported that malaria parasite infection has the highest prevalence among OIs in HIV infected children in Benin City Nigeria. This difference may be due to the fact that our study was conducted during the dry season while the environmental condition of the locality of the Benin study and the low immune status of the children may have resulted to a high prevalence of malaria infection.

This study also showed a significant association between PTB, Pneumonia, Oral Thrush and WHO clinical Staging. Place of residence and socioeconomic status were found to be significant risk factors for development of opportunistic infections. Of all opportunistic infections, 71% (23) occurred among rural dwellers. This may be due to the fact that rural dwellers have poor access to prompt health care as many specialized hospitals are located in big cities. Similarly, prevalence of opportunistic infections increased with worsening WHO stage because patients in this stage are severely immune suppressed. Prevalence of all the studied infections increased with lower social class. This may be due to the fact that indigent patients are unable to provide adequate nutrition, hence are unable to build the right immunity to fight infections.

Interpretation of results

This study depicts the fact that there is a high rate of opportunistic infections in HIV infected children. They prey on the immune suppression of HIV infected children and cause debilitating illness. Therefore physicians managing HIV infected children should look out for opportunistic infections especially in those with worsening WHO clinical staging, those from low socioeconomic background and those in rural areas. This is due to the fact that HIV infected children with these risk factors have very low immunity, malnutrition due to poverty and poor access to health care. Also, early diagnosis and prompt treatment leads to better outcomes. Furthermore, a multicenter study would give a broader view of opportunistic infections in HIV infected children.

Generalisability: This is a single center study, a multicenter study would have given the results a broader look.

Future research: Multi center study in patients with very low CD4 counts.

Limitations of study

This is a single center study. A multi center study may provide a broader view of OIs in HIV infected children.

Conclusion

In this study, the prevalence of OIs in HIV infected children at FMC Owerri remained high. OIs majorly occurred with worsening WHO clinical staging, low socioeconomic status and inhabitants of rural areas. We recommend consideration be given to early screening and treatment of OIs in HIV infected children.

What is already known on this topic

- Opportunistic infections are the most common cause of death among children living with HIV/AIDS.
- Opportunistic infections take advantage of the weakened immune system and cause devastating illnesses.
- Studies with respect to the prevalence of opportunistic infections in HIV infected Nigerian children are very limited.

What study adds

- The prevalence of OIs in HIV infected children at FUTH Owerri is high.
- OIs majorly occurred with worsening WHO clinical staging.
- OIs majorly occurred with low socioeconomic status and inhabitants of rural areas.

Declarations

Conflict of interest

Authors declare no conflict of interest.

UN's declaration of human right

All authors agree with UN's declaration of human right.

Author contributions

Chinelo Vivian Okeke: conceptualization of study, writing, expertise, review, data collection, data analysis, result interpretation.

Sampson Dayo Ejikunle: writing, review, result interpretation, expertise

Bede Ikenna Nnolim: writing, review, result interpretation, data analysis.

Idabo Oghenetega: writing, review, data analysis

Ganiyat Yusuf: writing, review, data analysis

Maria-lauretta Orji: conceptualization, supervision, writing, expertise, review.

Acknowledgements

The authors acknowledge the resident doctors in pediatrics department FUTHO and laboratory staff of HAART clinic FUTHO.

Funding

The authors funded the study and publication.

References

- [1] AVERT: Global HIV and AIDS epidemic. 2012. Available from www.avert.org/global-hiv-aids-epidemic.htm.
- [2] UNAIDS: Progress report on the Global plan towards the elimination of new HIV infection among children by 2015 and keeping their mothers alive. 2013. Available from www.unaids.org/en/media/unaids/contentassets/documents/unaids/publication/2013/20130625 Progress plan.en.pdf p63
- [3] National Agency for the control of Aids NACA: Nigeria Prevalence Rate 2019. [https://naca.gov.ng/Nigeria-prevalence rate.](https://naca.gov.ng/Nigeria-prevalence-rate)
- [4] Tindyebwa D, Kayita J, Musoke P, Eley B, Nduati R, Tumwesigye N et al, editors. HIV virology, pathogenesis and natural history. In; Handbook on paediatrics AIDS in Africa. African Network for the care of children affected by AIDS. 2nd ed. Uganda: Fam Healt Int. 2011;17-26.
- [5] Pediatric OI guidelines; Available from http://aidsinfo.nih.gov/contentfiles/Tvguidelines/oi_guidelines_pediatricspdf.
- [6] The Ethiopian Federal Ministry of Health. National ART guidelines for prevention, care and treatment. Ministry of health Ethiopia 2017.
- [7] Alarcon JO, Freimanis-Hance L, Krauss M, Reyes MF, Cardoso CA, Mussi-Pinhata MM, et al. Opportunistic and other infections in HIV infected children in Latin America compared to a similar cohort in the United States. AIDS Res Hum Retrivir. 2012; 28:282-8.
- [8] Gona P, Van Dyke RB, Williams PL, Dankner WM, Chernoff MC, Nachman SA, Seage GR. Incidence of opportunistic and other infections in HIV- Infected children in the HAART era. Jama. 2006; 296:292-300.
- [9] Modi S, Chui A, Ngeno B, Kellerman SE, Sugandhi N, Muhe L. Understanding the contribution of common childhood illnesses and opportunistic infections to morbidity and mortality in children living with HIV in resource limited settings. Aids. 2013;27:S159-67.
- [10] Ravichandra KR, Bikash RP and Sunil A. Opportunistic infections in HIV infected children and its correlation with CD4 count. Int J ContempPediatr. 2017; 4:1743-7.
- [11] Imade PE and Eghafona NO. Prevalent Opportunistic Infections associated with HIV -Positive children 0-5 years in Benin city Nigeria. Malaysian Journal of Microbiology. 2008;11-14.
- [12] Legal notice on publication of the details of the breakdown of the national and state provisional total 2006 census. Constitution of Federal Republic of Nigeria 2007;94:175
- [13] 2017 PHC priority tables- National Population Commission. Population.gov.ng.
- [14] Melkamu MW, Gebeyehu MT, Afenigus A et al. Incidence of common opportunistic infections among HIV- infected children on ART at Debre Markos referral hospital, Northwest Ethiopia a retrospective cohort study. BMC Infec Dis 2020; 20:50.
- [15] Prasitsuebsai W, Kariminia A, Puthanakit T, Lumbiganon P, Hansudewechakul R, Moy FS, et al. Impact of antiretroviral therapy on opportunistic infections of HIV-

infected children in the TREAT Asia pediatric HIV observational database. *PediatrInfec Dis J.* 2014; 33:747.

- [16] Ylitalo N, Brogly S, Hughes MD, Nachman S, Dankner W, VanDyke R et al. Risk factors for opportunistic illnesses in children with HIV in the era of HAART. *Arch PediatrAdolesc Med.* 2006; 160:778-87.



Published by AMMS Journal, this is an Open Access article distributed under the terms of the Creative Commons Attribution 4.0 International License. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2025